

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTALAB1643

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS 1 Web Page for STN Seminar Schedule - N. America  
NEWS 2 DEC 01 ChemPort single article sales feature unavailable  
NEWS 3 JUN 01 CAS REGISTRY Source of Registration (SR) searching  
enhanced on STN  
NEWS 4 JUN 26 NUTRACEUT and PHARMAML no longer updated  
NEWS 5 JUN 29 IMSCOPROFILE now reloaded monthly  
NEWS 6 JUN 29 EPFULL adds Simultaneous Left and Right Truncation  
(SLART) to AB, MCLM, and TI fields  
NEWS 7 JUL 09 PATDPAFULL adds Simultaneous Left and Right  
Truncation (SLART) to AB, CLM, MCLM, and TI fields  
NEWS 8 JUL 14 USGENE enhances coverage of patent sequence location  
(PSL) data  
NEWS 9 JUL 27 CA/CAPLUS enhanced with new citing references  
NEWS 10 JUL 16 GBFULL adds patent backfile data to 1855  
NEWS 11 JUL 21 USGENE adds bibliographic and sequence information  
NEWS 12 JUL 28 EPFULL adds first-page images and applicant-cited  
references  
NEWS 13 JUL 28 INPADOCDB and INPAFAMDB add Russian legal status data  
NEWS 14 AUG 08 Improve STN by completing a survey and be entered to  
win a gift card  
NEWS 15 AUG 10 Time limit for inactive STN sessions doubles to 40  
minutes  
NEWS 16 AUG 17 CAS REGISTRY, the Global Standard for Chemical  
Research, Approaches 50 Millionth Registration  
Milestone  
NEWS 17 AUG 18 COMPENDEX indexing changed for the Corporate Source  
(CS) field  
  
NEWS EXPRESS MAY 26 09 CURRENT WINDOWS VERSION IS V8.4,  
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use for software development or design, implementation of commercial gateways, or use of CAS and STN data in the building of commercial products is prohibited and may result in loss of user privileges and other penalties.

\*\*\*\*\*  
\* \*  
\* Please take a couple of minutes to complete our short survey. Your \*  
\* name will be entered to win one of five \$20 Amazon.com gift cards. \*  
\* \*  
\* See NEWS 14 for details or go directly to the survey at: \*  
\* <http://www.zoomerang.com/Survey/?p=WEB229H4S8Q5UL> \*  
\* \*  
\*\*\*\*\*

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 16:14:52 ON 18 AUG 2009

=> file caplus

| COST IN U.S. DOLLARS | ENTRY | SINCE FILE SESSION | TOTAL |
|----------------------|-------|--------------------|-------|
| FULL ESTIMATED COST  |       | 0.22               | 0.22  |

FILE 'CAPLUS' ENTERED AT 16:15:06 ON 18 AUG 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is

strictly prohibited.

FILE COVERS 1907 - 18 Aug 2009 VOL 151 ISS 8

FILE LAST UPDATED: 17 Aug 2009 (20090817/ED)

REVISED CLASS FIELDS (/NCL) LAST RELOADED: Jun 2009

USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Jun 2009

CAPLUS now includes complete International Patent Classification (IPC)  
reclassification data for the second quarter of 2009.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate  
substance identification.

The ALL, BIB, MAX, and STD display formats in the CA/CAPLUS family  
of databases have been updated to include new citing references  
information. This enhancement may impact record import into  
database management software. For additional information, refer  
to NEWS 9.

=> s (glypican-3 or GPC3 or DGSX or GTR2-2 or MXR7 or OCI5 or SDYS or SGB or  
SGBS or SGBS1) and melanoma

731 GLYPICAN

352 GLYPICANS

791 GLYPICAN

(GLYPICAN OR GLYPICANS)

7705968 3

310 GLYPICAN-3

(GLYPICAN(W)3)

283 GPC3

0 DGSX

24 GTR2

10178478 2

3 GTR2-2

(GTR2(W)2)

11 MXR7

7 OCI5

1 SDYS

251 SGB

62 SGBS

307 SGB

(SGB OR SGBS)

62 SGBS

3 SGBS1  
42855 MELANOMA  
4189 MELANOMAS  
19 MELANOMATA  
43425 MELANOMA  
(MELANOMA OR MELANOMAS OR MELANOMATA)  
L1 50 (GLYPICAN-3 OR GPC3 OR DGSX OR GTR2-2 OR MXR7 OR OCI5 OR  
SDYS  
OR SGB OR SGBS OR SGBS1) AND MELANOMA

=> duplicate remove L1

PROCESSING COMPLETED FOR L1

L2 50 DUPLICATE REMOVE L1 (0 DUPLICATES REMOVED)

=> s L2 (S) (diagnos? or assay? or measur? or detect?)

L3 50 S L2

PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH  
FIELD CODE - 'AND' OPERATOR ASSUMED 'L3 (S) '

363581 DIAGNOS?

655862 ASSAY?

3406766 MEASUR?

2007938 DETECT?

L4 30 L3 (S) (DIAGNOS? OR ASSAY? OR MEASUR? OR DETECT?)

=> s (glypican-3 or GPC3 or DGSX or GTR2-2 or MXR7 or OCI5 or SDYS or SGB or  
SGBS or SGBS1) (S) melanoma

731 GLYPICAN

352 GLYPICANS

791 GLYPICAN

(GLYPICAN OR GLYPICANS)

7705968 3

310 GLYPICAN-3

(GLYPICAN(W)3)

283 GPC3

0 DGSX

24 GTR2

10178478 2

3 GTR2-2

(GTR2(W)2)

11 MXR7

7 OCI5

1 SDYS

251 SGB

62 SGBS

307 SGB

(SGB OR SGBS)

62 SGBS  
3 SGBS1  
42855 MELANOMA  
4189 MELANOMAS  
19 MELANOMATA  
43425 MELANOMA  
(MELANOMA OR MELANOMAS OR MELANOMATA)  
L5 20 (GLYPICAN-3 OR GPC3 OR DGSX OR GTR2-2 OR MXR7 OR OCI5 OR  
SDYS  
OR SGB OR SGBS OR SGBS1) (S) MELANOMA

=> duplicate remove L5

PROCESSING COMPLETED FOR L5

L6 20 DUPLICATE REMOVE L5 (0 DUPLICATES REMOVED)

=> s L6 (S) (DIAGNOS? OR ASSAY? OR MEASUR? OR DETECT?)

L7 20 S L6

363581 DIAGNOS?

655862 ASSAY?

3406766 MEASUR?

2007938 DETECT?

L8 9 L7 (S) (DIAGNOS? OR ASSAY? OR MEASUR? OR DETECT?)

=> d L8 bib abs 1-9

L8 ANSWER 1 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2009:216149 CAPLUS

DN 151:144580

TI Glypican-3: A Novel Diagnostic Marker for Hepatocellular Carcinoma and  
More

AU Kandil, Dina H.; Cooper, Kumarasen

CS Department of Pathology, University of Vermont, Burlington, VT, 05401, USA

SO Advances in Anatomic Pathology (2009), 16(2), 125-129

CODEN: AAPDCK; ISSN: 1072-4109

PB Lippincott Williams & Wilkins

DT Journal; General Review

LA English

AB A review. Glypican-3 (GPC3) is a heparan sulfate proteoglycan that plays an important role in cell growth and differentiation. GPC3 function is tissue dependent. In some tissues, GPC3 acts as a tumor suppressor gene, whereas in others, it acts as an oncofetal protein. Studies have shown that GPC3 is a reliable marker for hepatocellular carcinoma. The sensitivity and specificity exceeds both .alpha.-fetoprotein and hepatocyte-paraffin1. GPC3 immunohistochem. can aid in the differentiation of testicular germ cell tumors, being expressed in all yolk sac tumors but not in seminomas. GPC3 expression has also been

identified in some squamous cell carcinomas of the lung and clear cell carcinomas of the ovary. The role of GPC3 in melanomas is still controversial. This article reviews the current information on the application of GPC3 immunostaining in surgical pathol. and cytol.

RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2007:407710 CAPLUS

DN 146:375330

TI Cancer metastasis diagnosis method, and therapeutic drug

IN Oguchi, Masao; Ishii, Keisuke

PA Perseus Proteomics Inc., Japan

SO Jpn. Kokai Tokkyo Koho, 18pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

| PATENT NO.          | KIND | DATE     | APPLICATION NO. | DATE     |
|---------------------|------|----------|-----------------|----------|
| PI JP 2007093274    | A    | 20070412 | JP 2005-279913  | 20050927 |
| PRAI JP 2005-279913 |      | 20050927 |                 |          |

AB Provided is a method for diagnosing cancer selected from Ewing's sarcoma primary nest, Ewing's sarcoma metastasis tissue, melanoma metastasis tissue and hepatic carcinoma metastasis tissue. The diagnostic method is characterized in that it comprises detecting GPC3 protein in a test sample (e.g., blood, blood serum, blood plasma) by an immunoassay using an anti-GPC3 antibody. Also provided is a diagnostic agent or therapeutic drug for metastatic cancer selected from Ewing's sarcoma primary nest, Ewing's sarcoma metastasis tissue, melanoma metastasis tissue and hepatic carcinoma metastasis tissue, which is characterized in that it contains resp. a GPC3 protein detection reagent contg. an anti-GPC3 antibody, or an anti-GPC3 antibody.

L8 ANSWER 3 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2006:383728 CAPLUS

DN 144:431112

TI Anti-SPARC and anti-glypican-33 antibodies and test kits for diagnosis of hepatic cancer and melanoma

IN Nishimura, Yasuharu; Nakatsura, Tetsuya; Ikuta, Yoshiaki

PA Kumamoto University, Japan

SO PCT Int. Appl., 24 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 1

| PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|---|------|----------|-----------------|----------|
| PI WO 2006043362  | A1   | 20060427 | WO 2005-JP14567 | 20050809 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW |      |          |                 |          |
| RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  |      |          |                 |          |
| AU 2005297303   | A1   | 20060427 | AU 2005-297303  | 20050809 |
| EP 1813943  | A1   | 20070801 | EP 2005-770440  | 20050809 |
| R: DE, FR, GB   |      |          |                 |          |
| US 20090111095  | A1   | 20090430 | US 2008-577435  | 20081212 |
| PRAI JP 2004-303688   | A    | 20041019 |                 |          |
| WO 2005-JP14567   | W    | 20050809 |                 |          |

AB It is intended to find another tumor marker useful for the early diagnosis of melanoma and to provide, utilizing the same, a diagnostic kit for malignant melanoma and method of diagnosis therefor. There is provided a diagnostic kit for malignant melanoma, comprising an antibody against SPARC and an antibody against GPC3.

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 4 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2006:39388 CAPLUS

DN 144:228826

TI Melanoma antigen gene family D 1 protein as hepatocarcinoma marker and its application in cancer diagnosis

IN Wan, Dafang; Gu, Jianren; Yang, Shengli

PA Shanghai New World Gene Technology Development Co., Ltd., Peop. Rep. China

SO Faming Zhuanli Shenqing Gongkai Shuomingshu, 22 pp.

CODEN: CNXXEV

DT Patent

LA Chinese

FAN.CNT 1

| PATENT NO.            | KIND | DATE     | APPLICATION NO.  | DATE     |
|-----------------------|------|----------|------------------|----------|
| PI CN 1629637         | A    | 20050622 | CN 2003-10109398 | 20031215 |
| CN 1281962            | C    | 20061025 |                  |          |
| PRAI CN 2003-10109398 |      | 20031215 |                  |          |

AB This invention relates to melanoma antigen gene family D1 protein (MAGFD1) as hepatocarcinoma marker, test kit and protein chip contg. anti-MAGED1 specific antibody for diagnosing hepatocarcinoma. The protein chip can also contains antibodies against other antigens, such as pTEN, p21, p27, p73, p53, Rb1, APC, nm23, P16, MXR7, IGF-II, TGF.alpha., HGF-R, c-erbB-1, Ras, Raf, c-myc and c-ets-2. This invention also describes medicine contg. antagonist of MAGFD1 and pharmaceutically acceptable carriers.

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L8 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2005:1234417 CAPLUS

DN 144:251411

TI Highly Sensitive Detection of Melanoma at an Early Stage Based on the Increased Serum Secreted Protein Acidic and Rich in Cysteine and Glypican-3 Levels

AU Ikuta, Yoshiaki; Nakatsura, Tetsuya; Kageshita, Toshiro; Fukushima, Satoshi; Ito, Shosuke; Wakamatsu, Kazumasa; Baba, Hideo; Nishimura, Yasuharu

CS Department of Immunogenetics, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

SO Clinical Cancer Research (2005), 11(22), 8079-8088

CODEN: CCREF4; ISSN: 1078-0432

PB American Association for Cancer Research

DT Journal

LA English

AB PURPOSE: There are no available tumor markers detecting primary melanoma at an early stage. The identification of such serum markers would be of significant benefit for an early diagnosis of melanoma. We recently identified glypican-3 (GPC3) as a novel tumor marker but could diagnose only 40% of melanomas. Thereby, we focused out attention on secreted protein acidic and rich in cysteine (SPARC) overexpressed in melanoma as another candidate for tumor marker. Exptl. Design: Secreted SPARC protein was quantified using ELISA in the sera from 109 melanoma patients, five patients with large congenital melanocytic nevus, 61 age-matched healthy donors, and 13 disease-free patients after undergoing a surgical removal. We also quantified GPC3 and 5-S-cysteinyl-dopa in the same serum samples and compared these markers for their diagnostic value. RESULTS: The serum SPARC concns. in melanoma patients were greater than those in healthy donors (P = 0.001). When we fixed a cutoff value at the mean concn. plus



2 SD of the healthy donors, the serum SPARC was found to have increased in the sera of 36 of the 109 (33%) melanoma patients, whereas there were three (4.9%) false-pos. cases of 61 healthy donors. Surprisingly, 19 of 36 patients showing increased SPARC levels were in stages 0 to II. The serum SPARC level decreased under the cutoff level in 10 of 13 patients after surgical removal. Using SPARC and GPC3 in combination thus enabled us to diagnose 47 of 75 (66.2%) melanoma patients at an early stage (0-II). CONCLUSIONS: SPARC or its combination with GPC3 is thus considered a potentially useful tumor marker, esp. for melanoma at an early stage.

OSC.G 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)

RE.CNT 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2005:448336 CAPLUS

DN 143:170230

TI Usefulness of the novel oncofetal antigen glypican-3  
for diagnosis of hepatocellular carcinoma and melanoma

AU Nakatsura, Tetsuya; Nishimura, Yasuharu

CS Department of Immunogenetics, Graduate School of Medical Sciences,  
Kumamoto University, Kumamoto, Japan

SO BioDrugs (2005), 19(2), 71-77

CODEN: BIDRF4; ISSN: 1173-8804

PB Adis International Ltd.

DT Journal; General Review

LA English

AB A review. Discussed are: novel strategies for identification of tumor-assocd. antigens; identification and expression of glypican (GPC)-3 in hepatocellular carcinoma and melanoma; detection of GPC-3 in patients; and known biol. properties of GPC-3.

OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)

RE.CNT 59 THERE ARE 59 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 7 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2005:395015 CAPLUS

DN 142:426401

TI Diagnostic agent for malignant melanoma

IN Nishimura, Yasuharu; Nakatsura, Tetsuya

PA Kumamoto Technology & Industry Foundation, Japan

SO PCT Int. Appl., 26 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 1

| PATENT NO.  | KIND | DATE     | APPLICATION NO.  | DATE     |
|---|------|----------|------------------|----------|
| PI WO 2005039380  | A2   | 20050506 | WO 2004-JP16374  | 20041028 |
| WO 2005039380   | A3   | 20050630 |                  |          |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW |      |          |                  |          |
| RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  |      |          |                  |          |
| AU 2004283614   | A1   | 20050506 | AU 2004-283614   | 20041028 |
| EP 1684076  | A2   | 20060726 | EP 2004-793354   | 20041028 |
| EP 1684076  | B1   | 20081231 |                  |          |
| R: CH, DE, FR, GB, LI, NL   |      |          |                  |          |
| CN 1894587  | A    | 20070110 | CN 2004-80032204 | 20041028 |
| US 20080044818  | A1   | 20080221 | US 2007-577343   | 20070305 |
| PRAI JP 2003-368725   | A    | 20031029 |                  |          |
| WO 2004-JP16374   | W    | 20041028 |                  |          |

AB A novel and clin. useful diagnostic agent for malignant melanoma. There is provided a diagnostic agent for malignant melanoma, comprising an antibody against GPC3, or a primer or probe capable of detecting the expression of GPC3.

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 8 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2004:1150198 CAPLUS

DN 142:278355

TI Mouse homologue of a novel human oncofetal antigen, glypican-3, evokes T-cell-mediated tumor rejection without autoimmune reactions in mice

AU Nakatsura, Tetsuya; Komori, Hiroyuki; Kubo, Tatsuko; Yoshitake, Yoshihiro; Senju, Satoru; Katagiri, Toyomasa; Furukawa, Yoichi; Ogawa, Michio; Nakamura, Yusuke; Nishimura, Yasuharu

CS Departments of Immunogenetics, Kumamoto University, Kumamoto, Japan

SO Clinical Cancer Research (2004), 10(24), 8630-8640

CODEN: CCREF4; ISSN: 1078-0432

PB American Association for Cancer Research

DT Journal

LA English

AB The authors recently identified glypican-3 (GPC3) overexpressed specifically in human hepatocellular carcinoma, as based on cDNA microarray anal. of 23,040 genes, and the authors reported that GPC3 is a novel tumor marker for human hepatocellular carcinoma and melanoma. GPC3, expressed in almost all hepatocellular carcinomas and melanomas, but not in normal tissues except for placenta or fetal liver, is a candidate of ideal tumor antigen for immunotherapy. In this study, the authors attempted to identify a mouse GPC3 epitope for CTLs in BALB/c mice, and for this, the authors set up a preclin. study to investigate the usefulness of GPC3 as a target for cancer immunotherapy in vivo. The authors identified a mouse GPC3-derived and Kd- restricted CTL epitope peptide in BALB/c mice. Inoculation of this GPC3 peptide-specific CTL into s.c. Colon26 cancer cells transfected with mouse GPC3 gene (C26/GPC3) led to rejection of the tumor in vivo, and i.v. inoculation of these CTLs into sublethally irradiated mice markedly inhibited growth of an established s.c. tumor. Inoculation of bone marrow-derived dendritic cells pulsed with this peptide prevented the growth of s.c. and splenic C26/GPC3 accompanied with massive infiltration of CD8+ T cells into tumors. Evidence of autoimmune reactions was never obsd. in surviving mice that had rejected tumor cell challenges. The authors found the novel oncofetal protein GPC3 to be highly immunogenic in mice and elicited effective antitumor immunity with no evidence of autoimmunity. GPC3 is useful not only for diagnosis of hepatocellular carcinoma and melanoma but also for possible immunotherapy or prevention of these tumors.

OSC.G 17 THERE ARE 17 CAPLUS RECORDS THAT CITE THIS RECORD (17 CITINGS)

RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 9 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2004:827193 CAPLUS

DN 142:4152

TI Identification of glypican-3 as a novel tumor marker for melanoma

AU Nakatsura, Tetsuya; Kageshita, Toshiro; Ito, Shosuke; Wakamatsu, Kazumasa; Monji, Mikio; Ikuta, Yoshiaki; Senju, Satoru; Ono, Tomomichi; Nishimura, Yasuharu

CS Department of Immunogenetics, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

SO Clinical Cancer Research (2004), 10(19), 6612-6621

CODEN: CCREF4; ISSN: 1078-0432

PB American Association for Cancer Research

DT Journal

LA English

AB The authors reported recently the novel tumor marker glypican-3 (GPC3) for hepatocellular carcinoma. In the present study, the authors investigated the expression of GPC3 in human melanoma cell lines and tissues and asked whether GPC3 could be a novel tumor marker for melanoma. Expression of GPC3 mRNA and protein was investigated in human melanoma cell lines and tissues using reverse transcription-PCR and immunohistochem. anal. Secreted GPC3 protein was quantified using ELISA in culture supernatants of melanoma cell lines and in sera from 91 patients with melanoma and 28 disease-free patients after surgical removal of primary melanoma. All of the subjects were Japanese nationals. In >80% of melanoma and melanocytic nevus, there was evident expression of GPC3 mRNA and protein. Furthermore, GPC3 protein was evidenced in sera of 39.6% (36 of 91) of melanoma patients but not in sera from subjects with large congenital melanocytic nevus (0 of 5) and from healthy donors (0 of 60). Twenty-seven of 36 serum GPC3-pos. patients were neg. for both serum 5-S-cysteinyl-dopa and melanoma-inhibitory activity, well-known tumor markers for melanoma. The pos. rate of serum GPC3 (39.6%) was significantly higher than that of 5-S-cysteinyl-dopa (26.7%) and of melanoma-inhibitory activity (20.9%). Surprisingly, the authors detected serum GPC3 even in patients with stage 0 in situ melanoma. The pos. rate of serum GPC3 at stage 0, I, and II (44.4%, 40.0%, and 47.6%) was significantly higher than that of 5-S-cysteinyl-dopa (0.0%, 8.0%, and 10.0%). Also obsd. was the disappearance of GPC3 protein in sera from 11 patients after surgical removal of the melanoma. GPC3 is apparently a novel tumor marker useful for the diagnosis of melanoma, esp. in early stages of the disorder.

OSC.G 33 THERE ARE 33 CAPLUS RECORDS THAT CITE THIS RECORD (33 CITINGS)

RE.CNT 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT